Light-cone spreading of perturbations in the Heisenberg spin chain

Abhishek Dhar
International centre for theoretical sciences
TIFR, Bangalore

Avijit Das, Saurish Chakrabarty (ICTS)

Subhro Bhattacharjee, Anupam Kundu, Samriddhi Sankar Ray (ICTS)

Roderich Moessner (MPI, Dresden)
David Huse (Princeton)

Phys. Rev. Lett. **121**, 024101 (2018)

Non-equilibrium behaviour of isolated classical and quantum systems, SISSA, Trieste, September 10-13, 2018

Outline

- Chaos in classical systems.
- Propagation of chaos in many-body systems.
- The out-of-time-ordered-commutator (OTOC) as a probe of quantum chaos and of ballistic propagation of information.
- Numerical results for classical Heisenberg spin-chain — Das et al, Phys. Rev. Lett. **121**, 024101 (2018).
- Numerical results for classical oscillator chains (Harmonic, Toda, FPU, Phi-4) — AD (in preparation)

Chaos in classical systems

A butterfly fluttering its wings in Beijing causes a tornado in Kansas!!

Classical Chaos - Sensitivity to initial conditions

Consider a dynamical system with deterministic time-evolution

$$\frac{dz_x}{dt} = f_x(\mathbf{z}) \ , \quad x = 1, 2, \ldots, N \ .$$

Consider two different initial conditions

$$z^A(0) \ \text{and} \ \ z^B(0) = z^A(0) + \delta z(0) \ .$$

After time t let

$$\delta z(t) = z^B(t) - z^A(t)$$



Definition of chaos — at large t

$$\lim_{\delta z(0) \to 0} \frac{|\delta z(t)|}{|\delta z(0)|} \sim e^{\lambda_{\mathbf{z}} t}$$

with $\lambda > 0$ — LYAPUNOV EXPONENT

Propagation of chaos

Suppose that the initial perturbation is localized in space

$$\delta z_x(0) = \epsilon \delta_{x,0} \ .$$

We expect that the differences in phase-space variables at any point in space should eventually diverge with time. Thus

$$\lim_{\delta z_0(0) \to 0} \frac{|\delta z_x(t)|}{|\delta z_0(0)|} \sim e^{\lambda_{\mathbf{z}}(x,t)t} \ .$$

Some questions:

- How long does it take before the perturbation is felt at point x ?
- How does the spatio-temporal evolution of the perturbation take place?

Spread of perturbations in a Hamiltonian system

Consider a chain of coupled oscillators (e.g coupled pendula).

$z_x = (q_x,\ p_x)$ — (position, momentum) of x^{th} particle. $[x = -N/2, \ldots, N/2]$.

- Disturb system locally.
- The spread of the perturbation can be expressed as a Poisson bracket:

$$\frac{\partial p_x(t)}{\partial p_0(0)} = -\{p_x(t), q_0(0)\} \qquad \{A, B\} = \sum_x \frac{\partial A}{\partial q_x}\frac{\partial B}{\partial p_x} - \frac{\partial A}{\partial p_x}\frac{\partial B}{\partial q_x}$$

- Linear response $\langle \partial p_x(t)/\partial p_0(0)\rangle_{eq} = -\langle\{p_x(t), q_0(0)\}\rangle = \beta\langle p_x(t)p_0(0)\rangle$
  —- Equilibrium correlation functions
- Chaos and "non-linear response" — $\left\langle \left(\frac{\partial p_x(t)}{\partial p_0(0)}\right)^2 \right\rangle$

Chaos in quantum systems

- $z \to \psi$ — No chaos since linear dynamics.
- Usual characterization of quantum chaos: level-spacing statistics
- Maldacena,Shenker,Stanford — replace Poisson bracket by commutator. Hence look at

$$-\langle [q_x(t), p_0(0)]^2 \rangle / \hbar^2 \ ,$$

  $< ... >$ represents expectation over pure or thermal state.
- No long time divergence but perhaps the intermediate-time growth shows a exponential growth regime and one can extract a Lyapunov exponent.
- Ehrenfest time and scrambling time.
- Ballistic spread characterized by butterfly velocity, Lieb-Robinson velocity.

Classical version of the OTOC

Let $\{Q_x, P_x\} = \{\delta q_x, \delta p_x\}$ be the perturbation of an initial condition $(\mathbf{q}, \mathbf{p})$.
Time evolution is given by:

$$\dot{q_x} = p_x \ , \quad \dot{p_x} = f_x$$
$$\dot{Q_x} = P_x, \quad \dot{P_x} = \frac{\partial f_x}{\partial q_y} Q_y = M_{xy}(\mathbf{q}) Q_y \ .$$

Solve with initial condition $(\mathbf{q}, \mathbf{p})$ chosen from an equilibrium distribution and $Q_x = 0, \ P_x = \delta_{x,0} \ , x = 0, 1, \ldots, N-1$.
Compute

$$C(x,t) = -\langle \{p_x(t), q_0\} \rangle = \langle P_x(t) \rangle, ---- \text{Correlations}$$
$$D(x,t) = \langle \{p_x(t), q_0\}^2 \rangle = \langle P_x^2(t) \rangle ---- \text{OTOC}$$

— Chaos propagation in the Heisenberg spin chain with precession dynamics.

Classical Heisenberg spin chains



Heisenberg spins $\mathbf{S} = (S^x, S^y, S^z)$ — unit three-dimensional vectors.
Hamiltonian given by

$$H = -\sum_{\ell=1}^{N} \mathbf{S}_\ell . \mathbf{S}_{\ell+1} \ .$$

Equations of motion [Structure of usual Hamiltonian dynamics]

$$\dot{\mathbf{S}}_\ell = \mathbf{S}_\ell \times \mathbf{B}_\ell^{\text{eff}} = \{\mathbf{S}_\ell, H\} \ .$$

Linear response at high temperatures

Expect diffusive hydrodynamic equations.

Spreading of a spin perturbation $C_{ss}(x,t) = \langle \mathbf{S}_x(t).\mathbf{S}_0(0)\rangle_{eq}$.



Energy correlations are similar.

Chaos and ballistic propagation in the *XXX* Heisenberg chain at high temerature



- Consider finite perturbation $\delta \mathbf{S}_0(0) = \epsilon \hat{\mathbf{n}}$ and look at

$$D(x,t) = \langle [\delta \mathbf{S}_x(t)]^2 \rangle / 2 = 1 - \langle \mathbf{S}_x^A(t).\mathbf{S}_x^B(t) \rangle$$

.

- Clearly $D(x,t) \to 1$ at large times
- However in the limit $\epsilon \to 0$

$$\left\langle \left(\frac{\partial \mathbf{S}_x(t)}{\partial \mathbf{S}_0(0)} \right)^2 \right\rangle = \frac{D(x,t)}{\epsilon^2} \sim e^{2\lambda t}$$

Spreading of perturbation for single realization

Space-time evolution of localized perturbation.



Light-cone evolution of $D(x, t)$

System of $N = 2048$ spins, average over random initial conditions.



The signal decays exponentially with distance away from the light cone $x = v_b t$, with $v \approx 1.67$.

Evolution of $D(x,t)$

Plot of $D(x,t)$ profile at different times (non-linear dynamics with finite ϵ).



From numerics, the front behaves as $D(x,t) \sim \epsilon^2 \; e^{2\lambda t[1-(x/v_b t)^2]}$

Behaviour of $D(x,t)$

Growth of $D(x,t)$ profile at different positions – from linear and non-linear dynamics.



From numerics, the signal grows as $D(x,t) \sim \epsilon^2 \; e^{2\lambda t[1-(x/v_b t)^2]}$, with $\lambda \approx 0.5$, $v_b \approx 1.67$.

Behaviour of $D(x,t)$

Estimating the butterfly speed - At what time does the signal reach some threshold value D_o at a given point x.



$x^{1/3}$ broadening of the front.

Connection to KPZ surface growth

- Exponential growth within the light-cone (Butterfly effect)
- Ballistic front (Butterfly velocity)
- Saturation at long times (for finite perturbation)
- Broadening of growth front - possily related to KPZ
  — A. S. Pikovsky and J. Kurths (1994)

Consider linearized dynamics. Define —

$$h(x,t) = \lim_{\epsilon \to 0} \log[\delta \mathbf{S}^2(x,t)/2\epsilon^2]/2$$

Our observations:

$$h(x,t) = at - bx^2/t + t^{1/3}\eta$$

Numerically $a \approx \lambda_L$ and $b \approx \lambda_L / v_b^2$.
This suggests that $h(x,t)$ is like the KPZ height field.
Distribution of η — Tracy-Widom ?

Connections to KPZ

Distribution of height at $x = 0$ at different times.



Clear $t^{1/3}$-scaling but no signs yet of Tracy-Widom.

Conclusions

- Investigated the classical version of “Out-of-Time-Ordered-Correlator” in the Heisenberg spin chain.

  —Ballistic front propagation in a diffusive system, exponential growth, broadening of front and possible connections to KPZ universality.

- $D(x,t) \sim e^{2\mu(x/t)t}, \quad \mu(v) = \lambda[1-(v/v_b)^2]$ — velocity-dependent Lyapunov exponent [R. J. Deissler and K. Kaneko (1987), S. Lepri, A. Politi, and A. Torcini (1996)]
  Integrable systems: $\mu(v) \sim -(1-v/v_b)^{3/2}$. [vanishes within light cone]

- Possible connections to the KPZ surface growth model.

- What physical observable ? Information propagation, Loschmidt echo,...

- OTOC in oscillator chains.

Numerical results: evolution of $D(x, t)$

Harmonic chain



Toda chain







Numerical results: evolution of $D(x,t)$

Harmonic chain



Toda chain



FPU-chain



phi-4 chain



Numerical results: evolution of $D(x,t)$

Harmonic chain



Toda chain





Exponential growth is seen in FPU case.